

# Expanding brackets and simplifying expressions

## A LEVEL LINKS

**Scheme of work:** 1a. Algebraic expressions – basic algebraic manipulation, indices and surds

## Key points

- When you expand one set of brackets you must multiply everything inside the bracket by what is outside.
- When you expand two linear expressions, each with two terms of the form  $ax + b$ , where  $a \neq 0$  and  $b \neq 0$ , you create four terms. Two of these can usually be simplified by collecting like terms.

## Examples

**Example 1** Expand  $4(3x - 2)$

$4(3x - 2) = 12x - 8$	Multiply everything inside the bracket by the 4 outside the bracket
-----------------------	---

**Example 2** Expand and simplify  $3(x + 5) - 4(2x + 3)$

$3(x + 5) - 4(2x + 3)$ $= 3x + 15 - 8x - 12$ $= 3 - 5x$	<p><b>1</b> Expand each set of brackets separately by multiplying <math>(x + 5)</math> by 3 and <math>(2x + 3)</math> by <math>-4</math></p> <p><b>2</b> Simplify by collecting like terms:  <math>3x - 8x = -5x</math> and <math>15 - 12 = 3</math></p>
---	--

**Example 3** Expand and simplify  $(x + 3)(x + 2)$

$(x + 3)(x + 2)$ $= x(x + 2) + 3(x + 2)$ $= x^2 + 2x + 3x + 6$ $= x^2 + 5x + 6$	<p><b>1</b> Expand the brackets by multiplying <math>(x + 2)</math> by <math>x</math> and <math>(x + 2)</math> by 3</p> <p><b>2</b> Simplify by collecting like terms:  <math>2x + 3x = 5x</math></p>
---	---

**Example 4** Expand and simplify  $(x - 5)(2x + 3)$

$(x - 5)(2x + 3)$ $= x(2x + 3) - 5(2x + 3)$ $= 2x^2 + 3x - 10x - 15$ $= 2x^2 - 7x - 15$	<p><b>1</b> Expand the brackets by multiplying <math>(2x + 3)</math> by <math>x</math> and <math>(2x + 3)</math> by <math>-5</math></p> <p><b>2</b> Simplify by collecting like terms:  <math>3x - 10x = -7x</math></p>
---	---

## Practice

1 Expand.

**a**  $3(2x - 1)$

**c**  $-(3xy - 2y^2)$

**b**  $-2(5pq + 4q^2)$

2 Expand and simplify.

**a**  $7(3x + 5) + 6(2x - 8)$

**c**  $9(3s + 1) - 5(6s - 10)$

**b**  $8(5p - 2) - 3(4p + 9)$

**d**  $2(4x - 3) - (3x + 5)$

3 Expand.

**a**  $3x(4x + 8)$

**c**  $-2h(6h^2 + 11h - 5)$

**b**  $4k(5k^2 - 12)$

**d**  $-3s(4s^2 - 7s + 2)$

4 Expand and simplify.

**a**  $3(y^2 - 8) - 4(y^2 - 5)$

**c**  $4p(2p - 1) - 3p(5p - 2)$

**b**  $2x(x + 5) + 3x(x - 7)$

**d**  $3b(4b - 3) - b(6b - 9)$

5 Expand  $\frac{1}{2}(2y - 8)$

6 Expand and simplify.

**a**  $13 - 2(m + 7)$

**b**  $5p(p^2 + 6p) - 9p(2p - 3)$

7 The diagram shows a rectangle.

Write down an expression, in terms of  $x$ , for the area of the rectangle.

Show that the area of the rectangle can be written as  $21x^2 - 35x$

$3x - 5$



$7x$

8 Expand and simplify.

**a**  $(x + 4)(x + 5)$

**c**  $(x + 7)(x - 2)$

**e**  $(2x + 3)(x - 1)$

**g**  $(5x - 3)(2x - 5)$

**i**  $(3x + 4y)(5y + 6x)$

**k**  $(2x - 7)^2$

**b**  $(x + 7)(x + 3)$

**d**  $(x + 5)(x - 5)$

**f**  $(3x - 2)(2x + 1)$

**h**  $(3x - 2)(7 + 4x)$

**j**  $(x + 5)^2$

**l**  $(4x - 3y)^2$

## Extend

9 Expand and simplify  $(x + 3)^2 + (x - 4)^2$

10 Expand and simplify.

**a**  $\left(x + \frac{1}{x}\right)\left(x - \frac{2}{x}\right)$

**b**  $\left(x + \frac{1}{x}\right)^2$

### Watch out!

When multiplying (or dividing) positive and negative numbers, if the signs are the same the answer is '+'; if the signs are different the answer is '-'.

# Answers

**1 a**  $6x - 3$

**c**  $-3xy + 2y^2$

**b**  $-10pq - 8q^2$

**2 a**  $21x + 35 + 12x - 48 = 33x - 13$

**b**  $40p - 16 - 12p - 27 = 28p - 43$

**c**  $27s + 9 - 30s + 50 = -3s + 59 = 59 - 3s$

**d**  $8x - 6 - 3x - 5 = 5x - 11$

**3 a**  $12x^2 + 24x$

**c**  $10h - 12h^3 - 22h^2$

**b**  $20k^3 - 48k$

**d**  $21s^2 - 21s^3 - 6s$

**4 a**  $-y^2 - 4$

$$\mathbf{c} \quad 2p - 7p^2$$

**b**  $5x^2 - 11x$

**d**  $6b^2$

**5**  $y - 4$

**6 a**  $-1 - 2m$

**b**  $5p^3 + 12p^2 + 27p$

**7**  $7x(3x - 5) = 21x^2 - 35x$

**8 a**  $x^2 + 9x + 20$

**c**  $x^2 + 5x - 14$

**e**  $2x^2 + x - 3$

g  $10x^2 - 31x + 15$

**i**  $18x^2 + 39xy + 20y^2$

**k**  $4x^2 - 28x + 49$

**b**  $x^2 + 10x + 21$

**d**  $x^2 - 25$

**f**  $6x^2 - x - 2$

## h $12x^2 + 13x - 14$

**j**  $x^2 + 10x + 25$

$$\mathbf{1} \quad 16x^2 - 24xy + 9y^2$$

**9**  $2x^2 - 2x + 25$

**10 a**  $x^2 - 1 - \frac{2}{x^2}$

**b**  $x^2 + 2 + \frac{1}{x^2}$

# Surds and rationalising the denominator

## A LEVEL LINKS

**Scheme of work:** 1a. Algebraic expressions – basic algebraic manipulation, indices and surds

## Key points

- A surd is the square root of a number that is not a square number, for example  $\sqrt{2}$ ,  $\sqrt{3}$ ,  $\sqrt{5}$ , etc.
- Surds can be used to give the exact value for an answer.
- $\sqrt{ab} = \sqrt{a} \times \sqrt{b}$
- $\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}$
- To rationalise the denominator means to remove the surd from the denominator of a fraction.
- To rationalise  $\frac{a}{\sqrt{b}}$  you multiply the numerator and denominator by the surd  $\sqrt{b}$
- To rationalise  $\frac{a}{b + \sqrt{c}}$  you multiply the numerator and denominator by  $b - \sqrt{c}$

## Examples

**Example 1** Simplify  $\sqrt{50}$

$\begin{aligned}\sqrt{50} &= \sqrt{25 \times 2} \\ &= \sqrt{25} \times \sqrt{2} \\ &= 5 \times \sqrt{2} \\ &= 5\sqrt{2}\end{aligned}$	<ol style="list-style-type: none"> <li>Choose two numbers that are factors of 50. One of the factors must be a square number</li> <li>Use the rule <math>\sqrt{ab} = \sqrt{a} \times \sqrt{b}</math></li> <li>Use <math>\sqrt{25} = 5</math></li> </ol>
---	---

**Example 2** Simplify  $\sqrt{147} - 2\sqrt{12}$

$\begin{aligned}\sqrt{147} - 2\sqrt{12} \\ &= \sqrt{49 \times 3} - 2\sqrt{4 \times 3} \\ &= \sqrt{49} \times \sqrt{3} - 2\sqrt{4} \times \sqrt{3} \\ &= 7 \times \sqrt{3} - 2 \times 2 \times \sqrt{3} \\ &= 7\sqrt{3} - 4\sqrt{3} \\ &= 3\sqrt{3}\end{aligned}$	<ol style="list-style-type: none"> <li>Simplify <math>\sqrt{147}</math> and <math>2\sqrt{12}</math>. Choose two numbers that are factors of 147 and two numbers that are factors of 12. One of each pair of factors must be a square number</li> <li>Use the rule <math>\sqrt{ab} = \sqrt{a} \times \sqrt{b}</math></li> <li>Use <math>\sqrt{49} = 7</math> and <math>\sqrt{4} = 2</math></li> <li>Collect like terms</li> </ol>
--	--

**Example 3** Simplify  $(\sqrt{7} + \sqrt{2})(\sqrt{7} - \sqrt{2})$

$  \begin{aligned}  &(\sqrt{7} + \sqrt{2})(\sqrt{7} - \sqrt{2}) \\  &= \sqrt{49} - \sqrt{7}\sqrt{2} + \sqrt{2}\sqrt{7} - \sqrt{4} \\  &= 7 - 2 \\  &= 5  \end{aligned}  $	<p><b>1</b> Expand the brackets. A common mistake here is to write <math>(\sqrt{7})^2 = 49</math></p> <p><b>2</b> Collect like terms:  <math display="block">-\sqrt{7}\sqrt{2} + \sqrt{2}\sqrt{7} = -\sqrt{7}\sqrt{2} + \sqrt{7}\sqrt{2} = 0</math></p>
---	---

**Example 4** Rationalise  $\frac{1}{\sqrt{3}}$

$  \begin{aligned}  \frac{1}{\sqrt{3}} &= \frac{1}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} \\  &= \frac{1 \times \sqrt{3}}{\sqrt{9}} \\  &= \frac{\sqrt{3}}{3}  \end{aligned}  $	<p><b>1</b> Multiply the numerator and denominator by <math>\sqrt{3}</math></p> <p><b>2</b> Use <math>\sqrt{9} = 3</math></p>
---	---

**Example 5** Rationalise and simplify  $\frac{\sqrt{2}}{\sqrt{12}}$

$  \begin{aligned}  \frac{\sqrt{2}}{\sqrt{12}} &= \frac{\sqrt{2}}{\sqrt{12}} \times \frac{\sqrt{12}}{\sqrt{12}} \\  &= \frac{\sqrt{2} \times \sqrt{4 \times 3}}{12} \\  &= \frac{2\sqrt{2}\sqrt{3}}{12} \\  &= \frac{\sqrt{2}\sqrt{3}}{6}  \end{aligned}  $	<p><b>1</b> Multiply the numerator and denominator by <math>\sqrt{12}</math></p> <p><b>2</b> Simplify <math>\sqrt{12}</math> in the numerator. Choose two numbers that are factors of 12. One of the factors must be a square number</p> <p><b>3</b> Use the rule <math>\sqrt{ab} = \sqrt{a} \times \sqrt{b}</math></p> <p><b>4</b> Use <math>\sqrt{4} = 2</math></p> <p><b>5</b> Simplify the fraction:  <math>\frac{2}{12}</math> simplifies to <math>\frac{1}{6}</math></p>
---	--

**Example 6** Rationalise and simplify  $\frac{3}{2+\sqrt{5}}$

$\frac{3}{2+\sqrt{5}} = \frac{3}{2+\sqrt{5}} \times \frac{2-\sqrt{5}}{2-\sqrt{5}}$ $= \frac{3(2-\sqrt{5})}{(2+\sqrt{5})(2-\sqrt{5})}$ $= \frac{6-3\sqrt{5}}{4+2\sqrt{5}-2\sqrt{5}-5}$ $= \frac{6-3\sqrt{5}}{-1}$ $= 3\sqrt{5}-6$	<p><b>1</b> Multiply the numerator and denominator by <math>2-\sqrt{5}</math></p> <p><b>2</b> Expand the brackets</p> <p><b>3</b> Simplify the fraction</p> <p><b>4</b> Divide the numerator by <math>-1</math> Remember to change the sign of all terms when dividing by <math>-1</math></p>
--	---

## Practice

**1** Simplify.

**a**  $\sqrt{45}$

**c**  $\sqrt{48}$

**e**  $\sqrt{300}$

**g**  $\sqrt{72}$

**b**  $\sqrt{125}$

**d**  $\sqrt{175}$

**f**  $\sqrt{28}$

**h**  $\sqrt{162}$

### Hint

One of the two numbers you choose at the start must be a square number.

**2** Simplify.

**a**  $\sqrt{72} + \sqrt{162}$

**c**  $\sqrt{50} - \sqrt{8}$

**e**  $2\sqrt{28} + \sqrt{28}$

**b**  $\sqrt{45} - 2\sqrt{5}$

**d**  $\sqrt{75} - \sqrt{48}$

**f**  $2\sqrt{12} - \sqrt{12} + \sqrt{27}$

### Watch out!

Check you have chosen the highest square number at the start.

**3** Expand and simplify.

**a**  $(\sqrt{2} + \sqrt{3})(\sqrt{2} - \sqrt{3})$

**c**  $(4 - \sqrt{5})(\sqrt{45} + 2)$

**b**  $(3 + \sqrt{3})(5 - \sqrt{12})$

**d**  $(5 + \sqrt{2})(6 - \sqrt{8})$

**4** Rationalise and simplify, if possible.

**a**  $\frac{1}{\sqrt{5}}$

**b**  $\frac{1}{\sqrt{11}}$

**c**  $\frac{2}{\sqrt{7}}$

**d**  $\frac{2}{\sqrt{8}}$

**e**  $\frac{2}{\sqrt{2}}$

**f**  $\frac{5}{\sqrt{5}}$

**g**  $\frac{\sqrt{8}}{\sqrt{24}}$

**h**  $\frac{\sqrt{5}}{\sqrt{45}}$

**5** Rationalise and simplify.

**a**  $\frac{1}{3-\sqrt{5}}$

**b**  $\frac{2}{4+\sqrt{3}}$

**c**  $\frac{6}{5-\sqrt{2}}$

## Extend

**6** Expand and simplify  $(\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y})$

**7** Rationalise and simplify, if possible.

**a**  $\frac{1}{\sqrt{9}-\sqrt{8}}$

**b**  $\frac{1}{\sqrt{x}-\sqrt{y}}$

## Answers

1    **a**     $3\sqrt{5}$   
       **c**     $4\sqrt{3}$   
       **e**     $10\sqrt{3}$   
       **g**     $6\sqrt{2}$

**b**     $5\sqrt{5}$   
**d**     $5\sqrt{7}$   
**f**     $2\sqrt{7}$   
**h**     $9\sqrt{2}$

2    **a**     $15\sqrt{2}$   
       **c**     $3\sqrt{2}$   
       **e**     $6\sqrt{7}$

**b**     $\sqrt{5}$   
**d**     $\sqrt{3}$   
**f**     $5\sqrt{3}$

3    **a**     $-1$   
       **c**     $10\sqrt{5}-7$

**b**     $9-\sqrt{3}$   
**d**     $26-4\sqrt{2}$

4    **a**     $\frac{\sqrt{5}}{5}$   
       **c**     $\frac{2\sqrt{7}}{7}$   
       **e**     $\sqrt{2}$   
       **g**     $\frac{\sqrt{3}}{3}$

**b**     $\frac{\sqrt{11}}{11}$   
**d**     $\frac{\sqrt{2}}{2}$   
**f**     $\sqrt{5}$   
**h**     $\frac{1}{3}$

5    **a**     $\frac{3+\sqrt{5}}{4}$

**b**     $\frac{2(4-\sqrt{3})}{13}$

**c**     $\frac{6(5+\sqrt{2})}{23}$

6     $x-y$

7    **a**     $3+2\sqrt{2}$

**b**     $\frac{\sqrt{x}+\sqrt{y}}{x-y}$



# Rules of indices

## A LEVEL LINKS

**Scheme of work:** 1a. Algebraic expressions – basic algebraic manipulation, indices and surds

## Key points

- $a^m \times a^n = a^{m+n}$
- $\frac{a^m}{a^n} = a^{m-n}$
- $(a^m)^n = a^{mn}$
- $a^0 = 1$
- $a^{\frac{1}{n}} = \sqrt[n]{a}$  i.e. the  $n$ th root of  $a$
- $a^{\frac{m}{n}} = \sqrt[n]{a^m} = \left(\sqrt[n]{a}\right)^m$
- $a^{-m} = \frac{1}{a^m}$
- The square root of a number produces two solutions, e.g.  $\sqrt{16} = \pm 4$ .

## Examples

**Example 1** Evaluate  $10^0$

$10^0 = 1$	Any value raised to the power of zero is equal to 1
------------	---

**Example 2** Evaluate  $9^{\frac{1}{2}}$

$9^{\frac{1}{2}} = \sqrt{9}$ $= 3$	Use the rule $a^{\frac{1}{n}} = \sqrt[n]{a}$
------------------------------------	--

**Example 3** Evaluate  $27^{\frac{2}{3}}$

$27^{\frac{2}{3}} = \left(\sqrt[3]{27}\right)^2$ $= 3^2$ $= 9$	<p><b>1</b> Use the rule <math>a^{\frac{m}{n}} = \left(\sqrt[n]{a}\right)^m</math></p> <p><b>2</b> Use <math>\sqrt[3]{27} = 3</math></p>
--	--

**Example 4** Evaluate  $4^{-2}$

$4^{-2} = \frac{1}{4^2}$ $= \frac{1}{16}$	<p><b>1</b> Use the rule <math>a^{-m} = \frac{1}{a^m}</math></p> <p><b>2</b> Use <math>4^2 = 16</math></p>
---	--

**Example 5** Simplify  $\frac{6x^5}{2x^2}$

$\frac{6x^5}{2x^2} = 3x^3$	<p><math>6 \div 2 = 3</math> and use the rule <math>\frac{a^m}{a^n} = a^{m-n}</math> to give <math>\frac{x^5}{x^2} = x^{5-2} = x^3</math></p>
----------------------------	---

**Example 6** Simplify  $\frac{x^3 \times x^5}{x^4}$

$\frac{x^3 \times x^5}{x^4} = \frac{x^{3+5}}{x^4} = \frac{x^8}{x^4}$ $= x^{8-4} = x^4$	<p><b>1</b> Use the rule <math>a^m \times a^n = a^{m+n}</math></p> <p><b>2</b> Use the rule <math>\frac{a^m}{a^n} = a^{m-n}</math></p>
--	--

**Example 7** Write  $\frac{1}{3x}$  as a single power of  $x$

$\frac{1}{3x} = \frac{1}{3}x^{-1}$	<p>Use the rule <math>\frac{1}{a^m} = a^{-m}</math>, note that the fraction <math>\frac{1}{3}</math> remains unchanged</p>
------------------------------------	--

**Example 8** Write  $\frac{4}{\sqrt{x}}$  as a single power of  $x$

$\frac{4}{\sqrt{x}} = \frac{4}{x^{\frac{1}{2}}}$ $= 4x^{-\frac{1}{2}}$	<p><b>1</b> Use the rule <math>a^{\frac{1}{n}} = \sqrt[n]{a}</math></p> <p><b>2</b> Use the rule <math>\frac{1}{a^m} = a^{-m}</math></p>
--	--

## Practice

1 Evaluate.

**a**  $14^0$

**b**  $3^0$

**c**  $5^0$

**d**  $x^0$

2 Evaluate.

**a**  $49^{\frac{1}{2}}$

**b**  $64^{\frac{1}{3}}$

**c**  $125^{\frac{1}{3}}$

**d**  $16^{\frac{1}{4}}$

3 Evaluate.

**a**  $25^{\frac{3}{2}}$

**b**  $8^{\frac{5}{3}}$

**c**  $49^{\frac{3}{2}}$

**d**  $16^{\frac{3}{4}}$

4 Evaluate.

**a**  $5^{-2}$

**b**  $4^{-3}$

**c**  $2^{-5}$

**d**  $6^{-2}$

5 Simplify.

**a**  $\frac{3x^2 \times x^3}{2x^2}$

**b**  $\frac{10x^5}{2x^2 \times x}$

**c**  $\frac{3x \times 2x^3}{2x^3}$

**d**  $\frac{7x^3y^2}{14x^5y}$

**e**  $\frac{y^2}{y^{\frac{1}{2}} \times y}$

**f**  $\frac{c^{\frac{1}{2}}}{c^2 \times c^{\frac{3}{2}}}$

**g**  $\frac{(2x^2)^3}{4x^0}$

**h**  $\frac{x^{\frac{1}{2}} \times x^{\frac{3}{2}}}{x^{-2} \times x^3}$

### Watch out!

Remember that any value raised to the power of zero is 1. This is the rule  $a^0 = 1$ .

6 Evaluate.

**a**  $4^{-\frac{1}{2}}$

**b**  $27^{-\frac{2}{3}}$

**c**  $9^{-\frac{1}{2}} \times 2^3$

**d**  $16^{\frac{1}{4}} \times 2^{-3}$

**e**  $\left(\frac{9}{16}\right)^{-\frac{1}{2}}$

**f**  $\left(\frac{27}{64}\right)^{-\frac{2}{3}}$

7 Write the following as a single power of  $x$ .

**a**  $\frac{1}{x}$

**b**  $\frac{1}{x^7}$

**c**  $\sqrt[4]{x}$

**d**  $\sqrt[5]{x^2}$

**e**  $\frac{1}{\sqrt[3]{x}}$

**f**  $\frac{1}{\sqrt[3]{x^2}}$

8 Write the following without negative or fractional powers.

**a**  $x^{-3}$

**b**  $x^0$

**c**  $x^{\frac{1}{5}}$

**d**  $x^{\frac{2}{5}}$

**e**  $x^{-\frac{1}{2}}$

**f**  $x^{-\frac{3}{4}}$

9 Write the following in the form  $ax^n$ .

**a**  $5\sqrt{x}$

**b**  $\frac{2}{x^3}$

**c**  $\frac{1}{3x^4}$

**d**  $\frac{2}{\sqrt{x}}$

**e**  $\frac{4}{\sqrt[3]{x}}$

**f** 3

## Extend

10 Write as sums of powers of  $x$ .

**a**  $\frac{x^5 + 1}{x^2}$

**b**  $x^2 \left( x + \frac{1}{x} \right)$

**c**  $x^{-4} \left( x^2 + \frac{1}{x^3} \right)$

# Answers

**1 a** 1

**b** 1

**c** 1

**d** 1

**2 a** 7

**b** 4

**c** 5

**d** 2

**3 a** 125

**b** 32

**c** 343

**d** 8

**4 a**  $\frac{1}{25}$

**b**  $\frac{1}{64}$

**c**  $\frac{1}{32}$

**d**  $\frac{1}{36}$

**5 a**  $\frac{3x^3}{2}$

**b**  $5x^2$

**c**  $3x$

**d**  $\frac{y}{2x^2}$

**e**  $y^{\frac{1}{2}}$

**f**  $c^{-3}$

**g**  $2x^6$

**h**  $x$

**6 a**  $\frac{1}{2}$

**b**  $\frac{1}{9}$

**c**  $\frac{8}{3}$

**d**  $\frac{1}{4}$

**e**  $\frac{4}{3}$

**f**  $\frac{16}{9}$

**7 a**  $x^{-1}$

**b**  $x^{-7}$

**c**  $x^{\frac{1}{4}}$

**d**  $x^{\frac{2}{5}}$

**e**  $x^{-\frac{1}{3}}$

**f**  $x^{-\frac{2}{3}}$

**8 a**  $\frac{1}{x^3}$

**b** 1

**c**  $\sqrt[5]{x}$

**d**  $\sqrt[5]{x^2}$

**e**  $\frac{1}{\sqrt{x}}$

**f**  $\frac{1}{\sqrt[4]{x^3}}$

**9 a**  $5x^{\frac{1}{2}}$

**b**  $2x^{-3}$

**c**  $\frac{1}{3}x^{-4}$

**d**  $2x^{-\frac{1}{2}}$

**e**  $4x^{-\frac{1}{3}}$

**f**  $3x^0$

**10 a**  $x^3 + x^{-2}$

**b**  $x^3 + x$

**c**  $x^{-2} + x^{-7}$

# Factorising expressions

## A LEVEL LINKS

Scheme of work: 1b. Quadratic functions – factorising, solving, graphs and the discriminants

## Key points

- Factorising an expression is the opposite of expanding the brackets.
- A quadratic expression is in the form  $ax^2 + bx + c$ , where  $a \neq 0$ .
- To factorise a quadratic equation find two numbers whose sum is  $b$  and whose product is  $ac$ .
- An expression in the form  $x^2 - y^2$  is called the difference of two squares. It factorises to  $(x - y)(x + y)$ .

## Examples

**Example 1** Factorise  $15x^2y^3 + 9x^4y$

$$15x^2y^3 + 9x^4y = 3x^2y(5y^2 + 3x^2)$$

The highest common factor is  $3x^2y$ .  
So take  $3x^2y$  outside the brackets and then divide each term by  $3x^2y$  to find the terms in the brackets

**Example 2** Factorise  $4x^2 - 25y^2$

$$4x^2 - 25y^2 = (2x + 5y)(2x - 5y)$$

This is the difference of two squares as the two terms can be written as  $(2x)^2$  and  $(5y)^2$

**Example 3** Factorise  $x^2 + 3x - 10$

$$b = 3, ac = -10$$

$$\text{So } x^2 + 3x - 10 = x^2 + 5x - 2x - 10$$

$$= x(x + 5) - 2(x + 5)$$

$$= (x + 5)(x - 2)$$

- 1 Work out the two factors of  $ac = -10$  which add to give  $b = 3$  (5 and -2)
- 2 Rewrite the  $b$  term ( $3x$ ) using these two factors
- 3 Factorise the first two terms and the last two terms
- 4  $(x + 5)$  is a factor of both terms

**Example 4** Factorise  $6x^2 - 11x - 10$

$b = -11, ac = -60$  So $6x^2 - 11x - 10 = 6x^2 - 15x + 4x - 10$ $= 3x(2x - 5) + 2(2x - 5)$ $= (2x - 5)(3x + 2)$	<ol style="list-style-type: none"> <li>1 Work out the two factors of <math>ac = -60</math> which add to give <math>b = -11</math> (-15 and 4)</li> <li>2 Rewrite the <math>b</math> term (<math>-11x</math>) using these two factors</li> <li>3 Factorise the first two terms and the last two terms</li> <li>4 <math>(2x - 5)</math> is a factor of both terms</li> </ol>
---	--

**Example 5** Simplify  $\frac{x^2 - 4x - 21}{2x^2 + 9x + 9}$

$\frac{x^2 - 4x - 21}{2x^2 + 9x + 9}$  For the numerator: $b = -4, ac = -21$  So $x^2 - 4x - 21 = x^2 - 7x + 3x - 21$ $= x(x - 7) + 3(x - 7)$ $= (x - 7)(x + 3)$  For the denominator: $b = 9, ac = 18$  So $2x^2 + 9x + 9 = 2x^2 + 6x + 3x + 9$ $= 2x(x + 3) + 3(x + 3)$ $= (x + 3)(2x + 3)$  So $\frac{x^2 - 4x - 21}{2x^2 + 9x + 9} = \frac{(x - 7)(x + 3)}{(x + 3)(2x + 3)}$ $= \frac{x - 7}{2x + 3}$	<ol style="list-style-type: none"> <li>1 Factorise the numerator and the denominator</li> <li>2 Work out the two factors of <math>ac = -21</math> which add to give <math>b = -4</math> (-7 and 3)</li> <li>3 Rewrite the <math>b</math> term (<math>-4x</math>) using these two factors</li> <li>4 Factorise the first two terms and the last two terms</li> <li>5 <math>(x - 7)</math> is a factor of both terms</li> <li>6 Work out the two factors of <math>ac = 18</math> which add to give <math>b = 9</math> (6 and 3)</li> <li>7 Rewrite the <math>b</math> term (<math>9x</math>) using these two factors</li> <li>8 Factorise the first two terms and the last two terms</li> <li>9 <math>(x + 3)</math> is a factor of both terms</li> <li>10 <math>(x + 3)</math> is a factor of both the numerator and denominator so cancels out as a value divided by itself is 1</li> </ol>
---	---

## Practice

1 Factorise.

a  $6x^4y^3 - 10x^3y^4$

c  $25x^2y^2 - 10x^3y^2 + 15x^2y^3$

b  $21a^3b^5 + 35a^5b^2$

2 Factorise

a  $x^2 + 7x + 12$

c  $x^2 - 11x + 30$

e  $x^2 - 7x - 18$

g  $x^2 - 3x - 40$

b  $x^2 + 5x - 14$

d  $x^2 - 5x - 24$

f  $x^2 + x - 20$

h  $x^2 + 3x - 28$

3 Factorise

a  $36x^2 - 49y^2$

c  $18a^2 - 200b^2c^2$

b  $4x^2 - 81y^2$

4 Factorise

a  $2x^2 + x - 3$

c  $2x^2 + 7x + 3$

e  $10x^2 + 21x + 9$

b  $6x^2 + 17x + 5$

d  $9x^2 - 15x + 4$

f  $12x^2 - 38x + 20$

5 Simplify the algebraic fractions.

a  $\frac{2x^2 + 4x}{x^2 - x}$

c  $\frac{x^2 - 2x - 8}{x^2 - 4x}$

e  $\frac{x^2 - x - 12}{x^2 - 4x}$

b  $\frac{x^2 + 3x}{x^2 + 2x - 3}$

d  $\frac{x^2 - 5x}{x^2 - 25}$

f  $\frac{2x^2 + 14x}{2x^2 + 4x - 70}$

6 Simplify

a  $\frac{9x^2 - 16}{3x^2 + 17x - 28}$

c  $\frac{4 - 25x^2}{10x^2 - 11x - 6}$

b  $\frac{2x^2 - 7x - 15}{3x^2 - 17x + 10}$

d  $\frac{6x^2 - x - 1}{2x^2 + 7x - 4}$

### Hint

Take the highest common factor outside the bracket.

## Extend

7 Simplify  $\sqrt{x^2 + 10x + 25}$

8 Simplify  $\frac{(x+2)^2 + 3(x+2)^2}{x^2 - 4}$



## Answers

- |          |                                    |                                 |
|----------|------------------------------------|---------------------------------|
| <b>1</b> | <b>a</b> $2x^3y^3(3x - 5y)$        | <b>b</b> $7a^3b^2(3b^3 + 5a^2)$ |
|          | <b>c</b> $5x^2y^2(5 - 2x + 3y)$    |                                 |
| <b>2</b> | <b>a</b> $(x + 3)(x + 4)$          | <b>b</b> $(x + 7)(x - 2)$       |
|          | <b>c</b> $(x - 5)(x - 6)$          | <b>d</b> $(x - 8)(x + 3)$       |
|          | <b>e</b> $(x - 9)(x + 2)$          | <b>f</b> $(x + 5)(x - 4)$       |
|          | <b>g</b> $(x - 8)(x + 5)$          | <b>h</b> $(x + 7)(x - 4)$       |
| <b>3</b> | <b>a</b> $(6x - 7y)(6x + 7y)$      | <b>b</b> $(2x - 9y)(2x + 9y)$   |
|          | <b>c</b> $2(3a - 10bc)(3a + 10bc)$ |                                 |
| <b>4</b> | <b>a</b> $(x - 1)(2x + 3)$         | <b>b</b> $(3x + 1)(2x + 5)$     |
|          | <b>c</b> $(2x + 1)(x + 3)$         | <b>d</b> $(3x - 1)(3x - 4)$     |
|          | <b>e</b> $(5x + 3)(2x + 3)$        | <b>f</b> $2(3x - 2)(2x - 5)$    |
| <b>5</b> | <b>a</b> $\frac{2(x+2)}{x-1}$      | <b>b</b> $\frac{x}{x-1}$        |
|          | <b>c</b> $\frac{x+2}{x}$           | <b>d</b> $\frac{x}{x+5}$        |
|          | <b>e</b> $\frac{x+3}{x}$           | <b>f</b> $\frac{x}{x-5}$        |
| <b>6</b> | <b>a</b> $\frac{3x+4}{x+7}$        | <b>b</b> $\frac{2x+3}{3x-2}$    |
|          | <b>c</b> $\frac{2-5x}{2x-3}$       | <b>d</b> $\frac{3x+1}{x+4}$     |
| <b>7</b> | $(x + 5)$                          |                                 |
| <b>8</b> | $\frac{4(x+2)}{x-2}$               |                                 |